THC DOSING GUIDE

MEDICATION:

THC : CBD RATIO:


INDICATION:

MG/ML:

SCHEDULE: NOCTE / PRN (AS REQUIRED)

STARTING DOSE: 1 drop / 0.05mL

Gradually increase dose until symptoms are improved or controlled. Stop or decrease if side effects occur. Do not take more than the maximum written on the prescription label.



IMPORTANT: There are 20 drops per mL. 1 drop = 0.05mL.

Max = 100mg/day or as directed by your prescriber.

Remember

DAY	1	2	3	4	5	6	7
BED	1x 0.05mL	2x O.1mL	3x () 0.15mL	4x O.2mL	5x () 0.25mL	6x O.3mL	7 x () 0.35mL
PRN (as required)	IF REQUIRED 0.05mL	IF REQUIRED 0.1mL	IF REQUIRED 0.15mL	IF REQUIRED 0.2mL	IF REQUIRED 0.25mL	IF REQUIRED 0.3mL	IF REQUIRED 0.35mL

Administration Instructions

- 1. Start by eating something containing fat (i.e. a spoonful of yoghurt, cheese, coconut oil, avocado).
- 2. Place drop/mL(s) in mouth and swirl for 90 seconds, then swallow.
- **3.** Reassess if desired symptom control has been achieved, noting THC can take up to 2 hours for full effect. If the answer's no, continue to titrate as per dosing table.
- 4. Once desired symptom control has been achieved, you've found your dosing regimen. Maintain your minimum effective dose.
- 5. Record your optimal dose in dosing diary.
- 1. Start Low and Go Slow.
- 2. When your symptoms have been controlled, STOP increasing your dose. Maintain this effective dose as required.
- 3. The goal is to find the minimum effective dose.